


Hanakāpī'ai, Kaua'i



WATERSHED FEATURES

Hanakāpī'ai watershed occurs on the island of Kaua'i. The Hawaiian meaning of the name is "bay sprinkling food". The area of the watershed is 3.8 square mi (9.8 square km), with maximum elevation of 4222 ft (1287 m). The watershed's DAR cluster code is 2, meaning that the watershed is small, steep in the upper watershed, and with little embayment. The percent of the watershed in the different land use districts is as follows: 0% agricultural, 100% conservation, 0% rural, and 0% urban.

Land Stewardship: Percentage of the land in the watershed managed or controlled by the corresponding agency or entity. Note that this is not necessarily ownership.

<u>Military</u>	<u>Federal</u>	<u>State</u>	<u>OHA</u>	<u>County</u>	<u>Nature Conservancy</u>	<u>Other Private</u>
0.0	0.0	100.0	0.0	0.0	0.0	0.0

Land Management Status: Percentage of the watershed in the categories of biodiversity protection and management created by the Hawaii GAP program.

Permanent Biodiversity <u>Protection</u>	Managed for Multiple <u>Uses</u>	Protected but <u>Unmanaged</u>	<u>Unprotected</u>
0.0	59.1	40.9	0.0

Land Use: Areas of the various categories of land use. These data are based on NOAA C-CAP remote sensing project.

	<u>Percent</u>	<u>Square mi</u>	<u>Square km</u>
High Intensity Developed	0.0	0.00	0.00
Low Intensity Developed	0.0	0.00	0.00
Cultivated	0.0	0.00	0.00
Grassland	0.7	0.03	0.07
Scrub/Shrub	80.8	3.07	7.95
Evergreen Forest	18.1	0.69	1.79
Palustrine Forested	0.0	0.00	0.00
Palustrine Scrub/Shrub	0.0	0.00	0.00
Palustrine Emergent	0.0	0.00	0.00
Estuarine Forested	0.0	0.00	0.00
Bare Land	0.3	0.01	0.03
Unconsolidated Shoreline	0.0	0.00	0.00
Water	0.0	0.00	0.00
Unclassified	0.0	0.00	0.00

STREAM FEATURES

Hanakāpī'ai is a perennial stream. Total stream length is 16.2 mi (26.1 km). The terminal stream order is 3.

Reach Type Percentages: The percentage of the stream's channel length in each of the reach type categories.

<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
0.0	0.8	20.2	26.6	52.4

The following stream(s) occur in the watershed:
Hanakāpī'ai

BIOTIC SAMPLING EFFORT

Biotic samples were gathered in the following year(s):

1965	1977	1979	1980	1984	1986	1989
1990	1992	1993	1994	1995	1999	2003

Distribution of Biotic Sampling: The number of survey locations that were sampled in the various reach types.

<u>Survey type</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
Damselfly Surveys	0	4	2	2	0
DAR Point Quadrat	0	50	202	0	0
HDFG	0	0	2	0	0
Microhabitat Survey	0	12	28	0	0
Published Report	0	3	5	1	0
Unpublished Report	0	0	5	1	0
USGS Surveys	0	0	1	0	0

BIOTA INFORMATION**Species List****Native Species**

Crustaceans	<i>Atyoida bisulcata</i>
	<i>Macrobrachium grandimanus</i>
Fish	<i>Awaous guamensis</i>
	<i>Eleotris sandwicensis</i>
	Gobiid sp.
	<i>Kuhlia sandwicensis</i>
	<i>Kuhlia xenura</i>
	<i>Lentipes concolor</i>
	<i>Sicyopterus stimpsoni</i>
	<i>Stenogobius hawaiiensis</i>
Snails	<i>Ferrissia sharpi</i>
	<i>Neritina granosa</i>
Worms	<i>Oligochaete sp.</i>

Native Species

Insects	Empidid sp.
	Ephydrid sp.
	<i>Limonia sp.</i>
	<i>Megalagrion eudytum</i>
	<i>Megalagrion heterogamias</i>
	<i>Megalagrion oresitrophum</i>
	<i>Megalagrion sp.</i>
	<i>Megalagrion vagabundum</i>
	<i>Orthocladus sp.</i>
	<i>Procanace sp.</i>
	<i>Thambemyia acrosticalis</i>
	Tipulid sp.

Introduced Species

Amphibians	<i>Bufo marinus</i>
	Ranid sp.
Crustaceans	<i>Macrobrachium lar</i>
	<i>Procambarus clarkii</i>
Fish	<i>Poecilia reticulata</i>
Snails	Physid sp.
	Planorbid sp.
Worms	<i>Southwellina hispida</i>

Introduced Species

Insects	<i>Cheumatopsyche analis</i>
	Chironomid larvae
	<i>Ischnura posita</i>
	<i>Ischnura ramburi</i>
	Trichoptera larvae

Species Size Data: Species size (inches) observed in DAR Point Quadrat Surveys.

<u>Scientific Name</u>	<u>Status</u>	<u>Minimum Size</u>	<u>Maximum Size</u>	<u>Average Size</u>
<i>Bufo marinus</i>	Introduced	1	2.5	1.8
<i>Atyoida bisulcata</i>	Endemic	1	1.5	1.3
<i>Macrobrachium grandimanus</i>	Endemic	0.75	0.75	0.8
<i>Macrobrachium lar</i>	Introduced	1	6	3.5

<i>Eleotris sandwicensis</i>	Endemic	2	6.5	4.3
<i>Kuhlia xenura</i>	Endemic	0.75	4	1.7
<i>Lentipes concolor</i>	Endemic	0.63	5	2.7
<i>Sicyopterus stimpsoni</i>	Endemic	0.5	25	2.7
<i>Awaous guamensis</i>	Indigenous	0.75	12	3.7
<i>Neritina granosa</i>	Endemic	0.25	2	0.9

Average Density: The densities (#/square yard) for species observed in DAR Point Quadrat Surveys averaged over all sample dates in each reach type.

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Low</u>	<u>Mid</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Atyoida bisulcata</i>	Endemic			0.05		
<i>Eleotris sandwicensis</i>	Endemic		0.17			
<i>Kuhlia xenura</i>	Endemic		2.53			
<i>Lentipes concolor</i>	Endemic		0.35	3.94		
<i>Macrobrachium grandimanus</i>	Endemic			0.01		
<i>Neritina granosa</i>	Endemic		0.39	0.72		
<i>Sicyopterus stimpsoni</i>	Endemic		15.8	9.3		
<i>Awaous guamensis</i>	Indigenous		2.62	1.05		
<i>Bufo marinus</i>	Introduced			0.03		
<i>Macrobrachium lar</i>	Introduced		0.35	0.31		

Species Distributions: Presence (P) of species in different stream reaches.

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Atyoida bisulcata</i>	Endemic			P	P	
<i>Macrobrachium grandimanus</i>	Endemic			P		
<i>Eleotris sandwicensis</i>	Endemic		P	P		
<i>Kuhlia xenura</i>	Endemic		P	P		
<i>Lentipes concolor</i>	Endemic		P	P	P	
<i>Sicyopterus stimpsoni</i>	Endemic		P	P	P	
<i>Megalagrion eudytum</i>	Endemic				P	
<i>Megalagrion heterogamias</i>	Endemic			P		
<i>Megalagrion oresitrophum</i>	Endemic		P			
<i>Megalagrion sp.</i>	Endemic			P		
<i>Megalagrion vagabundum</i>	Endemic		P		P	
<i>Orthocladus sp.</i>	Endemic			P		
<i>Thambemyia acrosticalis</i>	Endemic			P		
<i>Ferrissia sharpi</i>	Endemic			P		
<i>Neritina granosa</i>	Endemic		P	P		
<i>Awaous guamensis</i>	Indigenous		P	P		
<i>Gobiid sp.</i>	Indigenous			P		
<i>Limonia sp.</i>	Indigenous			P		
<i>Procanace sp.</i>	Indigenous			P		
<i>Bufo marinus</i>	Introduced			P		

Ranid sp.	Introduced		P	
<i>Macrobrachium lar</i>	Introduced	P	P	
<i>Cheumatopsyche analis</i>	Introduced		P	P
Chironomid larvae	Introduced		P	P
<i>Ischnura posita</i>	Introduced		P	P
<i>Ischnura ramburi</i>	Introduced	P	P	
Trichoptera larvae	Introduced		P	
Physid sp.	Introduced		P	
Planorbid sp.	Introduced		P	
Empidid sp.	Undetermined		P	
Ephydrid sp.	Undetermined		P	
Tipulid sp.	Undetermined		P	P
<i>Oligochaete sp.</i>	Undetermined		P	

HISTORIC RANKINGS

Historic Rankings: These are rankings of streams from historical studies. "Yes" means the stream was considered worthy of protection by that method. Some methods include non-biotic data in their determination. See Atlas Key for details.

Multi-Attribute Prioritization of Streams - Potential Heritage Streams (1998): Yes

Hawaii Stream Assessment Rank (1990): Outstanding

U.S. Fish and Wildlife Service High Quality Stream (1988): Yes

The Nature Conservancy- Priority Aquatic Sites (1985): No

National Park Service - Nationwide Rivers Inventory (1982): Yes

Current DAR Decision Rule Status: The following criteria are used by DAR to consider the biotic importance of streams. "Yes" means that watershed has that quality.


Native Insect Diversity <u>> 19 spp.</u>	Native Macrofauna <u>Diversity > 5 spp.</u>	Absence of Priority 1 <u>Introduced</u>
No	Yes	No
Abundance of Any <u>Native Species</u>	Presence of Candidate <u>Endangered Species</u>	Endangered Newcomb's <u>Snail Habitat</u>
No	No	Yes

CURRENT WATERSHED AND STREAM RATINGS


The current watershed and stream ratings are based on the data contained in the DAR Aquatic Surveys Database. The ratings provide the score for the individual watershed or stream, the distribution of ratings for that island, and the distribution of ratings statewide. This allows a better understanding of the meaning of a particular ranking and how it compares to other streams. The ratings are standardized to range from 0 to 10 (0 is lowest and 10 is highest rating) for each variable and the totals are also standardized so that the rating is not the average of each component rating. These ratings are subject to change as more data are entered into the DAR Aquatic Surveys Database and can be automatically recalculated as the data improve. In addition to the ratings, we have also provided an estimate of the confidence level of the ratings. This is called rating strength. The higher the rating strength the more likely the data and rankings represent the actual condition of the watershed, stream, and aquatic biota.

WATERSHED RATING: Hanakāpī'ai, Kaua'i


Land Cover Rating: Rating is based on a scoring system where in general forested lands score positively and developed lands score negatively.



Shallow Waters Rating: Rating is based on a combination of the extent of estuarine and shallow marine areas associated with the watershed and stream.





Stewardship Rating: Rating is based on a scoring system where higher levels of land and biodiversity protection within the watershed score positively.



WATERSHED RATING (Cont): Hanakāpī'ai, Kaua'i



Size Rating: Rating is based on the watershed area and total stream length. Larger watersheds and streams score more positively.

Watershed
Rating
4





Wetness Rating: Rating is based on the average annual rainfall within the watershed. Higher rainfall totals score more positively.

Watershed
Rating
5





Reach Diversity Rating: Rating is based on the types and amounts of different stream reaches available in the watershed. More area in different reach types score more positively.

Watershed
Rating
6



Total Watershed Rating: Rating is based on combination of Land Cover Rating, Shallow Waters Rating, Stewardship Rating, Size Rating, Wetness Rating, and Reach Diversity Rating.



Watershed
Rating
7



BIOLOGICAL RATING: Hanakāpī'ai, Kaua'i



Native Species Rating: Rating is based on the number of native species observed in the watershed.

Stream
Rating
10





Introduced Genera Rating: Rating is based on the number of introduced genera observed in the watershed.

Stream
Rating
7





All Species' Score Rating: Rating is based on the Hawaii Stream Assessment scoring system where native species score positively and introduced species score negatively.

Stream
Rating
6



Total Biological Rating: Rating is the combination of the Native Species Rating, Introduced Genera Rating, and the All Species' Score Rating.

Stream
Rating
7




OVERALL RATING: Hanakāpī'ai, Kaua'i

Overall Rating: Rating is a combination of the Total Watershed Rating and the Total Biological Rating.

Watershed
Rating


8

**RATING STRENGTH: Hanakāpī'ai, Kaua'i**

Rating Strength: Represents an estimate of the overall study effort in the stream and is a combination of the number of studies, number of different reaches surveyed, and the number of different survey types.

Information
Rating

9

**REFERENCES**

1966. Shima, S.I. Limnological Survey for Introduction of Exotic Species of Fish.
1979. Ford, J.I. Biology of a Hawaiian Fluvial Gastropod *Neritina granosa* Sowerby (Prosobranchia: Neritidae). Masters Thesis.
1980. Parrish, J.D. Numerical equivalents of biological data in the Lentipes report.
1980. Timbol, A.S., Sutter, A.J. and J.D. Parrish. Distribution and Relative Abundance of the Endemic Freshwater Goby, *Lentipes concolor* in Hawaii. Hawaii Cooperative Fishery Research Unit.

1980. Timbol, A.S., Sutter, A.J. and J.D. Parrish. Distribution, Relative Abundance, and Stream Environment of *Lentipes concolor* (Gill, 1860), and Associated Fauna in Hawaiian Streams.
1984. Kinzie III, R.A, Ford, J.I., Yuen, A.R. and S.J.L. Chow. Habitat Utilization Curves for Native Hawaiian Stream Fishes. Special Report No. 6:20:84.
1986. Nishimoto, R.T. and J.M. Fitzsimons. Courtship, Territoriality, and Coloration in the Endemic Hawaiian Freshwater Goby, *Lentipes Concolor*. Reproductive Ecology of *Lentipes concolor*. 811-817.
1991. Baker, J.A. Sampling Hawaiian Stream Gobies. Proceedings of the 1990 Symposium on Freshwater Stream Biology and Fisheries Management. 238-278.
1991. Honigman, L. and A. Newman. A Biological Database of Aquatic Resources on Hawaiian Streams. Proceedings of the 1990 Symposium on Freshwater Stream Biology and Fisheries Management. 51-76.
1991. Zink, R.M. Genetic Variation Within and Between Populations of *Lentipes concolor* from Hawai'i and Kaua'i. Proceedings of the 1990 Symposium on Freshwater Stream Biology and Fisheries Management. 96-105.
1992. Nishimoto, R.T. Memo: Flood Threat. Hanakapi'ia Stream 1992..
1992. Tate, D.C., Kuamo'o, D.G. and W.F.M. Ishikawa. Recruitment of Larval Fishes into Kaua'i Streams after Hurricane Iniki.
1993. Kinzie III, R.A. Reproductive Biology of an Endemic, Amphidromous Goby, *Lentipes concolor* in Hawaiian Streams. Environmental Biology of Fishes, Vol. 37. 257-268.
1994. Fitzsimons, J.M. and R.T. Nishimoto. Use of Fish Behavior in Assessing the Effects of Hurricane Iniki on the Hawaiian island of Kaua'i. Environmental Biology of Fishes, Vol. 43. 39-50.
1994. Font, W.F. and D.C. Tate. Helminth Parasites of Native Hawaiian Freshwater Fishes: An Example of Extreme Ecological Isolation. The Journal of Parasitology, Vol. 80, No. 5. 682-688.
1996. Fitzsimons, J.M. and R.T. Nishimoto. Recovery of Three Kaua'i Streams from Hurricane Iniki and Implications for the Restoration and Regeneration of Freshwater Ecosystems in Hawai'i. Will Stream Restoration Benefit Freshwater, Estuarine, and Marine Fisheries? 69-75.

1996. Font, W.F., Tate, D.C. and D.W. Llewellyn. Colonization of Native Hawaiian Stream Fishes by Helminth Parasites. Will Stream Restoration Benefit Freshwater, Estuarine, and Marine Fisheries? 94-111.
1998. Greenfield, D.W., Suzumoto, A.Y. and C. Chong. *Vitraria clarescens*, a Junior Synonym of the Freshwater Hawaiian Goby *Sicyopterus stimpsoni* (Teleostei: Gobiidae). COPEIA, Vol. 98 No. 2. 501-503.
1999. Radtke, R.L., Townsend, D.W., Kinzie, III, R.A. and D. Fey. Two-Dimensional X-ray Mapping Otoliths, a High-Resolution Technique for Determining Amphidromy in the Tropical Goby *Lentipes concolor*. Journal of Experimental Biology and Ecology, Vol. 238. 21-27.
2001. Radtke, R.L., Kinzie, III, R.A. and D.J. Shafer. Temporal and Spatial Variation in Length of Larval Life and Size Settlement of the Hawaiian Amphidromous Goby *Lentipes concolor*. The Journal of Fish Biology, Vol. 59. 928-938.
2002. Parham, J.E. Microhabitat Surveys from Dissertation - Spatial Models of Hawaiian Streams and Stream Fishes.
2005. Englund, R.A., Laederich, H.M., Ghotaslou, A., Bollick, H. and A.A. Scott. Analysis of Kaua'i Aquatic Invertebrate Benthic Samples. Final Report, Hawaii Biological Survey.
2005. USGS. Stream Quality Indicators of Hawai'i.
2006. Polhemus, D.A. Maps of Damselfly Locations.
2006. Polhemus, D.A. Megalagrion Survey Notes in spreadsheet form.
2008. Hawai'i Division of Aquatic Resources. DAR Point Quadrat Survey Data from the DAR Aquatic Surveys Database.

Blank Page