
Hanamā'ulu, Kaua'i



WATERSHED FEATURES

Hanamā'ulu watershed occurs on the island of Kaua'i. The Hawaiian meaning of the name is "tired [as of walking] bay". The area of the watershed is 13.4 square mi (34.8 square km), with maximum elevation of 1148 ft (350 m). The watershed's DAR cluster code is 4, meaning that the watershed is medium size, steep in the upper watershed, and with embayment. The percent of the watershed in the different land use districts is as follows: 73.9% agricultural, 6.9% conservation, 0% rural, and 19.2% urban.

Land Stewardship: Percentage of the land in the watershed managed or controlled by the corresponding agency or entity. Note that this is not necessarily ownership.

<u>Military</u>	<u>Federal</u>	<u>State</u>	<u>OHA</u>	County	Nature Conservancy	Other Private
0.0	0.1	4.5	0.0	0.0	0.0	95.4

Land Management Status: Percentage of the watershed in the categories of biodiversity protection and management created by the Hawaii GAP program.

Permanent Biodiversity	Managed for Multiple	Protected but	
<u>Protection</u>	<u>Uses</u>	<u>Unmanaged</u>	<u>Unprotected</u>
0.0	0.0	4.6	95.4

Land Use: Areas of the various categories of land use. These data are based on NOAA C-CAP remote sensing project.

	<u>Percent</u>	Square mi	Square km
High Intensity Developed	2.9	0.39	1.02
Low Intensity Developed	5.1	0.68	1.76
Cultivated	22.8	3.07	7.95
Grassland	15.1	2.03	5.27
Scrub/Shrub	22.4	3.01	7.79
Evergreen Forest	29.0	3.90	10.11
Palustrine Forested	0.0	0.00	0.00
Palustrine Scrub/Shrub	0.5	0.07	0.19
Palustrine Emergent	0.1	0.01	0.02
Estuarine Forested	0.0	0.00	0.00
Bare Land	0.8	0.11	0.29
Unconsolidated Shoreline	0.1	0.01	0.03
Water	1.2	0.16	0.40
Unclassified	0.0	0.00	0.00

STREAM FEATURES

Hanamā'ulu is a perennial stream. Total stream length is 30.9 mi (49.7 km). The terminal stream order is 3.

Reach Type Percentages: The percentage of the stream's channel length in each of the reach type categories.

Estuary	Lower	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
1.1	0.0	87.8	11.1	0.0

The following stream(s) occur in the watershed: Hanamā'ulu

BIOTIC SAMPLING EFFORT

Biotic samples were gathered in the following year(s): 1952 1962 1964 1978 1994 2003

Distribution of Biotic Sampling: The number of survey locations that were sampled in the various reach types.

Survey type	<u>Estuary</u>	Lower	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
Damselfly Surveys	0	0	4	0	0
HDFG	0	0	2	0	0
HDFG releases	0	0	1	0	0
Published Report	0	0	1	0	0
Reservoir	0	0	5	0	0

BIOTA INFORMATION

Species List

Native Species Native Species

Crustaceans Atyoida bisulcata Insects Anax junius Ostracod sp. Anax sp.

Megalagrion heterogamias

Megalagrion sp.

Megalagrion vagabundum

Introduced Species

Introduced Species Fish Clarias fuscus Insects Chironomid Iarvae

Gambusia affinis Enallagma civile Ischnura ramburi Lepomis macrochirus

Lepomis sp.

Micropterus dolomieu Micropterus salmoides

Micropterus sp. Poecilia reticulata

Tilapia sp.

unidentified cyprinid

Snails Melanoides tuberculata

Species found in Impoundments

Fish Clarias fuscus

> Lepomis sp. Micropterus sp. Tilapia sp.

unidentified cyprinid

Species Distributions: Presence (P) of species in different stream reaches.

Scientific Name	<u>Status</u>	<u>Estuary</u>	Lower	<u>Middle</u>	<u>Upper</u> <u>Headwaters</u>
Atyoida bisulcata	Endemic			Р	
Megalagrion heterogamias	Endemic			Р	
Megalagrion sp.	Endemic			Р	
Megalagrion vagabundum	Endemic			Р	

Anax junius	Indigenous	Р
Anax sp.	Indigenous	Р
Clarias fuscus	Introduced	Р
Gambusia affinis	Introduced	Р
Lepomis macrochirus	Introduced	Р
Lepomis sp.	Introduced	Р
Micropterus dolomieu	Introduced	Р
Micropterus salmoides	Introduced	Р
Micropterus sp.	Introduced	Р
Poecilia reticulata	Introduced	Р
Tilapia sp.	Introduced	Р
unidentified cyprinid	Introduced	Р
Chironomid larvae	Introduced	Р
Enallagma civile	Introduced	Р
Ischnura ramburi	Introduced	Р
Melanoides tuberculata	Introduced	Р
Ostracod sp.	Undetermined	Р

HISTORIC RANKINGS

Historic Rankings: These are rankings of streams from historical studies. "Yes" means the stream was considered worthy of protection by that method. Some methods include non-biotic data in their determination. See Atlas Key for details.

Multi-Attribute Prioritization of Streams - Potential Heritage Streams (1998): No

Hawaii Stream Assessment Rank (1990): Limited

U.S. Fish and Wildlife Service High Quality Stream (1988): No

The Nature Conservancy- Priority Aquatic Sites (1985): No

National Park Service - Nationwide Rivers Inventory (1982): No

Current DAR Decision Rule Status: The following criteria are used by DAR to consider the biotic importance of streams. "Yes" means that watershed has that quality.


Native Insect Diversity > 19 spp.	Native Macrotauna Diversity > 5 spp.	Absence of Priority 1 <u>Introduced</u>
No	No	No
Abundance of Any Native Species	Presence of Candidate Endangered Species	Endangered Newcomb's Snail Habitat
No	No	No

CURRENT WATERSHED AND STREAM RATINGS


The current watershed and stream ratings are based on the data contained in the DAR Aquatic Surveys Database. The ratings provide the score for the individual watershed or stream, the distribution of ratings for that island, and the distribution of ratings statewide. This allows a better understanding of the meaning of a particular ranking and how it compares to other streams. The ratings are standardized to range from 0 to 10 (0 is lowest and 10 is highest rating) for each variable and the totals are also standardized so that the rating is not the average of each component rating. These ratings are subject to change as more data are entered into the DAR Aquatic Surveys Database and can be automatically recalculated as the data improve. In addition to the ratings, we have also provided an estimate of the confidence level of the ratings. This is called rating strength. The higher the rating strength the more likely the data and rankings represent the actual condition of the watershed, stream, and aquatic biota.

WATERSHED RATING: Hanamā'ulu, Kaua'i


<u>Land Cover Rating</u>: Rating is based on a scoring sytem where in general forested lands score positively and developed lands score negatively.



<u>Shallow Waters Rating</u>: Rating is based on a combination of the extent of estuarine and shallow marine areas associated with the watershed and stream.





<u>Stewardship Rating</u>: Rating is based on a scoring system where higher levels of land and biodiversity protection within the watershed score positively.



WATERSHED RATING (Cont): Hanamā'ulu, Kaua'i

<u>Size Rating</u>: Rating is based on the watershed area and total stream length. Larger watersheds and streams score more positively.







Wetness Rating: Rating is based on the average annual rainfall within the watershed. Higher rainfall totals score more positively.

Watershed Rating

2







<u>Reach Diversity Rating</u>: Rating is based on the types and amounts of different stream reaches available in the watershed. More area in different reach types score more positively.

Watershed Rating

4







<u>Total Watershed Rating</u>: Rating is based on combination of <u>Land Cover Rating</u>, <u>Shallow Waters Rating</u>, <u>Stewardship Rating</u>, <u>Size Rating</u>, <u>Wetness Rating</u>, and <u>Reach Diversity Rating</u>.

Watershed Rating

4







BIOLOGICAL RATING: Hanamā'ulu, Kaua'i

<u>Native Species Rating</u>: Rating is based on the number of native species observed in the watershed.

Stream Rating

1







<u>Introduced Genera Rating</u>: Rating is based on the number of introduced genera observed in the watershed.

Stream Rating

5







<u>All Species' Score Rating:</u> Rating is based on the Hawaii Stream Assessment scoring system where native species score positively and introduced species score negatively.

Stream Rating

3







<u>Total Biological Rating</u>: Rating is the combination of the <u>Native Species Rating</u>, <u>Introduced Genera Rating</u>, and the <u>All Species' Score Rating</u>.

Stream Rating


2






OVERALL RATING: Hanamā'ulu, Kaua'i

Overall Rating: Rating is a combination of the <u>Total Watershed Rating</u> and the <u>Total Biological</u> Rating.



RATING STRENGTH: Hanamā'ulu, Kaua'i

<u>Rating Strength:</u> Represents an estimate of the overall study effort in the stream and is a combination of the number of studies, number of different reaches surveyed, and the number of different survey types.



REFERENCES

- 1965. Shima, S.I. Limnological Survey for Introduction of Exotic Species of Fish.
- 2005. Englund, R.A., Laederich, H.M., Ghotaslou, A., Bollick, H. and A.A. Scott. Analysis of Kaua'i Aquatic Invertebrate Benthic Samples. Final Report, Hawaii Biological Survey.
- 2006. Polhemus, D.A. Maps of Damselfly Locations.
- 2006. Polhemus, D.A. Megalagrion Survey Notes in spreadsheet form.
- 2008. Hawai'i Division of Aquatic Resources. DAR Point Quadrat Survey Data from the DAR Aquatic Surveys Database.
- 2008. Hawai'i Division of Aquatic Resources. Impoundment Surveys in DAR Aquatic Surveys Database.