


Nu'alolo, Kaua'i



WATERSHED FEATURES

Nu'alolo watershed occurs on the island of Kaua'i. The Hawaiian meaning of the name is unknown. The area of the watershed is 2.6 square mi (6.8 square km), with maximum elevation of 4190 ft (1277 m). The watershed's DAR cluster code is 2, meaning that the watershed is small, steep in the upper watershed, and with little embayment. The percent of the watershed in the different land use districts is as follows: 0% agricultural, 100% conservation, 0% rural, and 0% urban.

Land Stewardship: Percentage of the land in the watershed managed or controlled by the corresponding agency or entity. Note that this is not necessarily ownership.

<u>Military</u>	<u>Federal</u>	<u>State</u>	<u>OHA</u>	<u>County</u>	<u>Nature Conservancy</u>	<u>Other</u>	<u>Private</u>
0.0	0.0	100.0	0.0	0.0	0.0		0.0

Land Management Status: Percentage of the watershed in the categories of biodiversity protection and management created by the Hawaii GAP program.

Permanent Biodiversity <u>Protection</u>	Managed for Multiple <u>Uses</u>	Protected but <u>Unmanaged</u>	<u>Unprotected</u>
0.0	28.1	71.9	0.0

Land Use: Areas of the various categories of land use. These data are based on NOAA C-CAP remote sensing project.

	<u>Percent</u>	<u>Square mi</u>	<u>Square km</u>
High Intensity Developed	0.0	0.00	0.00
Low Intensity Developed	0.4	0.01	0.03
Cultivated	0.0	0.00	0.00
Grassland	0.2	0.01	0.02
Scrub/Shrub	63.9	1.67	4.33
Evergreen Forest	33.8	0.88	2.29
Palustrine Forested	0.0	0.00	0.00
Palustrine Scrub/Shrub	0.0	0.00	0.00
Palustrine Emergent	0.0	0.00	0.00
Estuarine Forested	0.0	0.00	0.00
Bare Land	1.5	0.04	0.10
Unconsolidated Shoreline	0.1	0.00	0.00
Water	0.1	0.00	0.01
Unclassified	0.0	0.00	0.00

STREAM FEATURES

Nu'alolo is a perennial stream. Total stream length is 6.6 mi (10.6 km). The terminal stream order is 2.

Reach Type Percentages: The percentage of the stream's channel length in each of the reach type categories.

<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
0.0	0.0	19.9	17.6	62.5

The following stream(s) occur in the watershed:
Nu'alolo

BIOTIC SAMPLING EFFORT

Biotic samples were gathered in the following year(s):

1979 1980 1984 1992 1993 1994

Distribution of Biotic Sampling: The number of survey locations that were sampled in the various reach types.

<u>Survey type</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
Damselfly Surveys	0	0	2	0	1
DAR Point Quadrat	0	9	167	2	0
Published Report	2	0	2	0	0

BIOTA INFORMATION**Species List****Native Species**

Crustaceans *Atyoida bisulcata*
Macrobrachium grandimanus

Fish *Awaous guamensis*
Eleotris sandwicensis
Lentipes concolor
Sicyopterus stimpsoni
Stenogobius hawaiiensis

Snails *Neritina granosa*

Worms *Southwellina hispida*

Native Species

Insects *Megalagrion oresitrophum*
Megalagrion vagabundum

Introduced Species

Amphibians *Bufo marinus*
Ranid sp.

Crustaceans *Macrobrachium lar*

Species Size Data: Species size (inches) observed in DAR Point Quadrat Surveys.

<u>Scientific Name</u>	<u>Status</u>	<u>Minimum Size</u>	<u>Maximum Size</u>	<u>Average Size</u>
Ranid sp.	Introduced	0.5	0.5	0.5
<i>Atyoida bisulcata</i>	Endemic	0.5	1.25	1.0
<i>Macrobrachium grandimanus</i>	Endemic	4	4	4.0
<i>Macrobrachium lar</i>	Introduced	1	6	3.5
<i>Eleotris sandwicensis</i>	Endemic	3	3	3.0
<i>Lentipes concolor</i>	Endemic	0.75	5	2.8
<i>Sicyopterus stimpsoni</i>	Endemic	0.75	6	3.1
<i>Awaous guamensis</i>	Indigenous	0.5	12	4.0
<i>Neritina granosa</i>	Endemic	0.5	2	0.9

Average Density: The densities (#/square yard) for species observed in DAR Point Quadrat Surveys averaged over all sample dates in each reach type.

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Low</u>	<u>Mid</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Atyoida bisulcata</i>	Endemic			0.03		
<i>Eleotris sandwicensis</i>	Endemic			0.03		

<i>Lentipes concolor</i>	Endemic		5.03
<i>Macrobrachium grandimanus</i>	Endemic		0.03
<i>Neritina granosa</i>	Endemic	6.83	0.36
<i>Sicyopterus stimpsoni</i>	Endemic	1.14	3.27
<i>Awaous guamensis</i>	Indigenous	2.28	1.16
<i>Macrobrachium lar</i>	Introduced	2.85	2.83
Ranid sp.	Introduced		0.15

Species Distributions: Presence (P) of species in different stream reaches.

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Atyoida bisulcata</i>	Endemic			P		
<i>Macrobrachium grandimanus</i>	Endemic			P		
<i>Eleotris sandwicensis</i>	Endemic			P		
<i>Lentipes concolor</i>	Endemic			P		
<i>Sicyopterus stimpsoni</i>	Endemic	P	P	P		
<i>Megalagrion oresitrophum</i>	Endemic			P		
<i>Megalagrion vagabundum</i>	Endemic			P		P
<i>Neritina granosa</i>	Endemic		P	P		
<i>Awaous guamensis</i>	Indigenous	P	P	P		
Ranid sp.	Introduced			P		
<i>Macrobrachium lar</i>	Introduced			P	P	

HISTORIC RANKINGS

Historic Rankings: These are rankings of streams from historical studies. "Yes" means the stream was considered worthy of protection by that method. Some methods include non-biotic data in their determination. See Atlas Key for details.

Multi-Attribute Prioritization of Streams - Potential Heritage Streams (1998): Yes

Hawaii Stream Assessment Rank (1990): Outstanding

U.S. Fish and Wildlife Service High Quality Stream (1988): Yes

The Nature Conservancy- Priority Aquatic Sites (1985): No

National Park Service - Nationwide Rivers Inventory (1982): Yes

Current DAR Decision Rule Status: The following criteria are used by DAR to consider the biotic importance of streams. "Yes" means that watershed has that quality.


Native Insect Diversity > 19 spp.	Native Macrofauna Diversity > 5 spp.	Absence of Priority 1 Introduced
No	Yes	No
Abundance of Any Native Species	Presence of Candidate Endangered Species	Endangered Newcomb's Snail Habitat
No	No	No

CURRENT WATERSHED AND STREAM RATINGS


The current watershed and stream ratings are based on the data contained in the DAR Aquatic Surveys Database. The ratings provide the score for the individual watershed or stream, the distribution of ratings for that island, and the distribution of ratings statewide. This allows a better understanding of the meaning of a particular ranking and how it compares to other streams. The ratings are standardized to range from 0 to 10 (0 is lowest and 10 is highest rating) for each variable and the totals are also standardized so that the rating is not the average of each component rating. These ratings are subject to change as more data are entered into the DAR Aquatic Surveys Database and can be automatically recalculated as the data improve. In addition to the ratings, we have also provided an estimate of the confidence level of the ratings. This is called rating strength. The higher the rating strength the more likely the data and rankings represent the actual condition of the watershed, stream, and aquatic biota.

WATERSHED RATING: Nu‘alolo, Kaua‘i


Land Cover Rating: Rating is based on a scoring system where in general forested lands score positively and developed lands score negatively.



Shallow Waters Rating: Rating is based on a combination of the extent of estuarine and shallow marine areas associated with the watershed and stream.




Stewardship Rating: Rating is based on a scoring system where higher levels of land and biodiversity protection within the watershed score positively.




WATERSHED RATING (Cont): Nu'alolo, Kaua'i


Size Rating: Rating is based on the watershed area and total stream length. Larger watersheds and streams score more positively.




Wetness Rating: Rating is based on the average annual rainfall within the watershed. Higher rainfall totals score more positively.



Reach Diversity Rating: Rating is based on the types and amounts of different stream reaches available in the watershed. More area in different reach types score more positively.





Total Watershed Rating: Rating is based on combination of Land Cover Rating, Shallow Waters Rating, Stewardship Rating, Size Rating, Wetness Rating, and Reach Diversity Rating.



BIOLOGICAL RATING: Nu'alolo, Kaua'i



Native Species Rating: Rating is based on the number of native species observed in the watershed.

Stream Rating
9





Introduced Genera Rating: Rating is based on the number of introduced genera observed in the watershed.

Stream Rating
8





All Species' Score Rating: Rating is based on the Hawaii Stream Assessment scoring system where native species score positively and introduced species score negatively.

Stream Rating
6




Total Biological Rating: Rating is the combination of the Native Species Rating, Introduced Genera Rating, and the All Species' Score Rating.

Stream Rating
7




OVERALL RATING: Nu‘alolo, Kaua‘i

Overall Rating: Rating is a combination of the Total Watershed Rating and the Total Biological Rating.



RATING STRENGTH: Nu‘alolo, Kaua‘i

Rating Strength: Represents an estimate of the overall study effort in the stream and is a combination of the number of studies, number of different reaches surveyed, and the number of different survey types.



REFERENCES

Unknown. Species Summary Report.

1980. Timbol, A.S., Sutter, A.J. and J.D. Parrish. Distribution and Relative Abundance of the Endemic Freshwater Goby, *Lentipes concolor* in Hawaii. Hawaii Cooperative Fishery Research Unit.

1980. Timbol, A.S., Sutter, A.J. and J.D. Parrish. Distribution, Relative Abundance, and Stream Environment of *Lentipes concolor* (Gill, 1860), and Associated Fauna in Hawaiian Streams.

1991. Honigman, L. and A. Newman. A Biological Database of Aquatic Resources on Hawaiian Streams. Proceedings of the 1990 Symposium on Freshwater Stream Biology and Fisheries Management. 51-76.

1994. Fitzsimons, J.M. and R.T. Nishimoto. Use of Fish Behavior in Assessing the Effects of Hurricane Iniki on the Hawaiian island of Kaua‘i. *Environmental Biology of Fishes*, Vol. 43. 39-50.
1994. Font, W.F. and D.C. Tate. Helminth Parasites of Native Hawaiian Freshwater Fishes: An Example of Extreme Ecological Isolation. *The Journal of Parasitology*, Vol. 80, No. 5. 682-688.
1996. Fitzsimons, J.M. and R.T. Nishimoto. Recovery of Three Kaua‘i Streams from Hurricane Iniki and Implications for the Restoration and Regeneration of Freshwater Ecosystems in Hawai‘i. *Will Stream Restoration Benefit Freshwater, Estuarine, and Marine Fisheries?* 69-75.
1996. Font, W.F., Tate, D.C. and D.W. Llewellyn. Colonization of Native Hawaiian Stream Fishes by Helminth Parasites. *Will Stream Restoration Benefit Freshwater, Estuarine, and Marine Fisheries?* 94-111.
1996. Tate, D.C. Effect of Larval and Postlarval Fish Behavior in Determining the Instream Distribution of Adult *Awaous guamensis* and *Lentipes concolor* in Hakalau Stream, Hawai‘i. *Will Stream Restoration Benefit Freshwater, Estuarine, and Marine Fisheries?* 132-147.
1997. Tate, D.C. The Role of Behavioral Interactions of Immature Hawaiian Stream Fishes (Pisces: Gobiodei) in Population Dispersal and Distribution. *Micronesia* (30) 1. 51-70.
2006. Polhemus, D.A. Megalagrion Survey Notes in spreadsheet form.
2008. Hawai‘i Division of Aquatic Resources. DAR Point Quadrat Survey Data from the DAR Aquatic Surveys Database.

Blank Page