
DAR Watershed Code: 33012

Moanalua, Oʻahu



WATERSHED FEATURES

Moanalua watershed occurs on the island of O'ahu. The Hawaiian meaning of the name is unknown. The area of the watershed is 10.5 square mi (27.2 square km), with maximum elevation of 2812 ft (857 m). The watershed's DAR cluster code is not yet determined. The percent of the watershed in the different land use districts is as follows: 0% agricultural, 65.3% conservation, 0% rural, and 34.7% urban.

Land Stewardship: Percentage of the land in the watershed managed or controlled by the corresponding agency or entity. Note that this is not necessarily ownership.

<u>Military</u>	Federal	<u>State</u>	<u>OHA</u>	<u>County</u>	Nature Conservancy	Other Private
0.0	17.8	4.6	0.0	43.6	0.0	34.0

Land Management Status: Percentage of the watershed in the categories of biodiversity protection and management created by the Hawaii GAP program.

Permanent Biodiversity	Managed for Multiple	Protected but	
Protection	Uses	<u>Unmanaged</u>	Unprotected
0.0	0.0	66.0	34.0

Land Use: Areas of the various categories of land use. These data are based on NOAA C-CAP remote sensing project.

	Percent	<u>Square mi</u>	<u>Square km</u>
High Intensity Developed	9.8	1.03	2.67
Low Intensity Developed	9.9	1.04	2.70
Cultivated	0.0	0.00	0.00
Grassland	5.2	0.54	1.41
Scrub/Shrub	47.5	4.99	12.93
Evergreen Forest	26.9	2.83	7.32
Palustrine Forested	0.0	0.00	0.00
Palustrine Scrub/Shrub	0.0	0.00	0.00
Palustrine Emergent	0.0	0.00	0.00
Estuarine Forested	0.0	0.00	0.01
Bare Land	0.4	0.04	0.11
Unconsolidated Shoreline	0.0	0.00	0.00
Water	0.3	0.03	0.08
Unclassified	0.0	0.00	0.00

STREAM FEATURES

Moanalua is a perennial stream. Total stream length is 24.5 mi (39.5 km). The terminal stream order is 3.

Reach Type Percentages: The percentage of the stream's channel length in each of the reach type categories.

<u>Estuary</u>	Lower	Middle	<u>Upper</u>	Headwaters
4.9	0.0	50.4	44.7	0.0

The following stream(s) occur in the watershed: Kahauiki Manaiki Moanalua

BIOTIC SAMPLING EFFORT

Biotic sa	mples were	gathered in t	he following	g year(s):		
1976	1990	1994	1995	1997	2000	2002
2006						

Distribution of Biotic Sampling: The number of survey locations that were sampled in the various reach types.

Survey type	<u>Estuary</u>	Lower	<u>Middle</u>	<u>Upper</u>	Headwaters
Damselfly Surveys	0	1	3	0	0
DAR Rapid BioAssessment	0	0	1	0	0
Published Report	0	2	3	1	0

Species List

BIOTA INFORMATION

Native Species	S	Native Species			
Crustaceans	Macrobrachium grandimanus Thalamita crenata	Insects	Anax junius Megalagrion xanthomelas		
Fish Snails	Awaous guamensis Eleotris sandwicensis Sphyraena barracuda Stenogobius hawaiiensis Neritina vespertina				
Introduced Sp	ecies	Introduced Species			
Crustaceans Fish	Macrobrachium lar Procambarus clarkii Clarias fuscus Gambusia affinis Poecilia reticulata Poecilia sphenops Tilapia sp. Xiphophorus helleri Xiphophorus maculatus	Insects	Crocothemis servilia Hydroptila potosina Ischnura ramburi Notonecta indica Orthemis ferruginea Pantala flavescens Tramea abdominalis		
Snails	Melanoides tuberculata Thiara granifera				

Species Distributions: Presence (P) of species in different stream reaches.

Scientific Name	<u>Status</u>	<u>Estuary</u>	Lower	<u>Middle</u>	Upper Headwaters
Megalagrion xanthomelas	Endemic			Р	
Thalamita crenata	Indigenous		Р		
Sphyraena barracuda	Indigenous		Р		
Anax junius	Indigenous			Р	
Macrobrachium lar	Introduced				Р
Procambarus clarkii	Introduced			Р	
Gambusia affinis	Introduced		Р		Р
Poecilia reticulata	Introduced		Р		Р
Tilapia sp.	Introduced		Р		Р
Xiphophorus helleri	Introduced		Р		Р
Crocothemis servilia	Introduced			Р	

Moanalua, Oʻahu

Hydroptila potosina	Introduced		Р	
Ischnura ramburi	Introduced	Р		
Notonecta indica	Introduced		Р	
Orthemis ferruginea	Introduced		Р	
Pantala flavescens	Introduced		Р	
Tramea abdominalis	Introduced		Р	
Thiara granifera	Introduced	Р		Ρ

HISTORIC RANKINGS

Historic Rankings: These are rankings of streams from historical studies. "Yes" means the stream was considered worthy of protection by that method. Some methods include non-biotic data in their determination. See Atlas Key for details.

Multi-Attribute Prioritization of Streams - Potential Heritage Streams (1998): No Hawaii Stream Assessment Rank (1990): not ranked U.S. Fish and Wildlife Service High Quality Stream (1988): No The Nature Conservancy- Priority Aquatic Sites (1985): No National Park Service - Nationwide Rivers Inventory (1982): No

Current DAR Decision Rule Status: The following criteria are used by DAR to consider the biotic importance of streams. "Yes" means that watershed has that quality.


Native Insect Diversity	Native Macrofauna	Absence of Priority 1
> 19 spp.	<u>Diversity > 5 spp.</u>	Introduced
No	Yes	No
Abundance of Any	Presence of Candidate	Endangered Newcomb's
<u>Native Species</u>	Endangered Species	<u>Snail Habitat</u>
No	Yes	No

CURRENT WATERSHED AND STREAM RATINGS


The current watershed and stream ratings are based on the data contained in the DAR Aquatic Surveys Database. The ratings provide the score for the individual watershed or stream, the distribution of ratings for that island, and the distribution of ratings statewide. This allows a better understanding of the meaning of a particular ranking and how it compares to other streams. The ratings are standardized to range from 0 to 10 (0 is lowest and 10 is highest rating) for each variable and the totals are also standardized so that the rating is not the average of each component rating. These ratings are subject to change as more data are entered into the DAR Aquatic Surveys Database and can be automatically recalculated as the data improve. In addition to the ratings, we have also provided an estimate of the confidence level of the ratings. This is called rating strength. The higher the rating strength the more likely the data and rankings represent the actual condition of the watershed, stream, and aquatic biota.

WATERSHED RATING: Moanalua, O'ahu


Land Cover Rating: Rating is based on a scoring sytem where in general forested lands score positively and developed lands score negatively.



<u>Shallow Waters Rating</u>: Rating is based on a combination of the extent of estuarine and shallow marine areas associated with the watershed and stream.




<u>Stewardship Rating</u>: Rating is based on a scoring system where higher levels of land and biodiversity protection within the watershed score positively.




WATERSHED RATING (Cont): Moanalua, O'ahu


<u>Size Rating</u>: Rating is based on the watershed area and total stream length. Larger watersheds and streams score more positively.




Wetness Rating: Rating is based on the average annual rainfall within the watershed. Higher rainfall totals score more positively.



<u>Reach Diversity Rating</u>: Rating is based on the types and amounts of different stream reaches available in the watershed. More area in different reach types score more positively.




Total Watershed Rating: Rating is based on combination of Land Cover Rating, Shallow Waters Rating, Stewardship Rating, Size Rating, Wetness Rating, and Reach Diversity Rating.




BIOLOGICAL RATING: Moanalua, O'ahu


<u>Native Species Rating</u>: Rating is based on the number of native species observed in the watershed.




Introduced Genera Rating: Rating is based on the number of introduced genera observed in the watershed.



<u>All Species' Score Rating:</u> Rating is based on the Hawaii Stream Assessment scoring system where native species score positively and introduced species score negatively.




<u>Total Biological Rating</u>: Rating is the combination of the <u>Native Species Rating</u>, <u>Introduced</u> <u>Genera Rating</u>, and the <u>All Species' Score Rating</u>.




OVERALL RATING: Moanalua, O'ahu

Overall Rating: Rating is a combination of the <u>Total Watershed Rating</u> and the <u>Total Biological</u> <u>Rating</u>.



RATING STRENGTH: Moanalua, O'ahu

<u>Rating Strength</u>: Represents an estimate of the overall study effort in the stream and is a combination of the number of studies, number of different reaches surveyed, and the number of different survey types.



REFERENCES

1976. Nishimoto, M. 1976 storm drain Moanalua Stream, Oahu. Stream shocking data.

- 1996. Polhemus, D.A. The Orangeblack Hawaiian Damselfly, Megalagrion xanthomelas (Odonata: Coenagrionidae): Clarifying the Current Range of a Threatened Species. Bishop Museum Occasional Papers: No. 45. 30-53.
- 2000. Tagawa, A.W. Management of a Database on the Occurrence, Abundance and Distribution of Native Freshwater Species. Job Progress Report.
- 2001. Englund, R.A., Preston, D.J., Evenhuis, N.L., Cowie, R.H., Imada, C., Puttock, C. and K. Arakaki. Long-term Monitoring of One the Most Restricted Insect Populations in the United States, Megalagrion xanthomelas ,Selys-Longchamps, at Tripler Army Medical Center, Oahu, Hawaii ,Zygoptera: Coenagrionidae. Odonatologica Vol. 30, No. 3. 255-263.

- 2002. Kido, M.H. Biological and Habitat Assessment of Moanalua Stream, O'ahu. Technical Report, City and County of Honolulu.
- 2006. Polhemus, D.A. Maps of Damselfly Locations.
- 2006. Polhemus, D.A. Megalagrion Survey Notes in spreadsheet form.
- 2008. Hawai'i Division of Aquatic Resources. Rapid Assessment Surveys in DAR Aquatic Surveys Database.

Blank Page